

Quenching behaviour of a nonlocal parabolic MEMS equation

Kin Ming Hui

Institute of Mathematics, Academia Sinica,
Nankang, Taipei, 11529, Taiwan, R. O. C.

March 16, 2010

Abstract

Let $\Omega \subset \mathbb{R}^n$ be a C^2 bounded domain and $\chi > 0$, $\lambda > 0$, be constants. We obtain upper bounds for the quenching time of the solutions of the nonlocal parabolic MEMS equation $u_t = \Delta u + \lambda/(1-u)^2(1 + \chi \int_{\Omega} 1/(1-u) dx)^2$ in $\Omega \times (0, \infty)$, $u = 0$ on $\partial\Omega \times (0, \infty)$, $u(x, 0) = u_0$ in Ω , when λ is large. We will prove the compactness of the quenching set under a mild condition on u_0 . When $\Omega = B_R = \{x \in \mathbb{R}^n : |x| < R\}$ and u_0 is radially symmetric and monotone decreasing in $0 \leq r \leq R$, we prove that the point $x = 0$ is the only possible quenching set. When $\Omega = B_R$ and u_0 is radially symmetric which also satisfies some strict concavity assumption, we prove that for any $\beta \in (2, 3)$ the solution satisfies $1 - u(x, t) \geq C|x|^{\frac{2}{\beta}}$ for some constant $C > 0$ and that the solution u quenches in a finite time for any sufficiently large $\lambda > 0$. We also obtain the quenching time estimate in this case.

Key words: parabolic nonlocal MEMS equation, quenching time estimates, compactness, quenching set

Mathematics Subject Classification: Primary 35B40 Secondary 35B05, 35K50, 35K20

Micro-electromechanical systems (MEMS) devices are key components of many electronic devices including accelerometers for airbag deployment in cars, inkjet printer heads, and the device for the protection of hard disk, etc. It is therefore interesting to understand the mathematical modelling of the MEMS devices and study the various properties of such models. Interested readers can read the book, Modeling MEMS and NEMS [22], by J.A. Pelesko and D.H. Bernstein for the mathematical modeling and various applications of MEMS devices.

One model of MEMS [10], [19], [22], consists of a fixed membrane and a deformable membrane of the same shape which is coated with a thin dielectric material and held fixed at the boundary. When no voltage is applied to the membranes, the two membranes are parallel to each other with one at a fixed small distance on top of the other. When a voltage is applied to the membranes, the deformable membrane will deflect towards the fixed membrane. Recently there are a lot of study on the equations arising from such model of MEMS by N. Ghoussoub, Y. Guo, Z. Pan and M.J. Ward [3], [4], [5], [6], [9], [10], J.S. Guo, B. Hu and C.J. Wang [8], K.M. Hui [14], [15], [16], N.I. Kavallaris, T. Miyasita and T. Suzuki [17], F. Lin and Y. Yang [19], L. Ma, Z. Guo and J.C. Wei [11], [12], [20], G. Flores, G.A. Mercado, J.A. Pelesko and A.A. Triolo [2], [21], [23] etc.

As observed by N. Ghoussoub, Y. Guo, J.A. Pelesko and others [3], [22], the deflection of the deformable membrane from its equilibrium position is modeled by the following parabolic equation,

$$\begin{cases} \frac{\partial u}{\partial t} = \Delta u + \frac{\lambda f(x)}{(1-u)^2} & \text{in } \Omega \times (0, T) \\ u = 0 & \text{on } \partial\Omega \times (0, T) \\ u(x, 0) = u_0 & \text{in } \Omega \end{cases} \quad (P'_\lambda)$$

where $\lambda \geq 0$ is a constant and $f \not\equiv 0$ is a nonnegative function on $\overline{\Omega}$ which depends on the dielectric constant of the coating on the membrane. When the voltage between the membranes are due to circuit series capacitance, the deflection of the deformable membrane from its equilibrium position is modeled by the following nonlocal parabolic equation [7],

$$\begin{cases} \frac{\partial u}{\partial t} = \Delta u + \frac{\lambda}{(1-u)^2(1 + \chi \int_{\Omega} \frac{dy}{1-u(y,t)})^2} & \text{in } \Omega \times (0, T) \\ u = 0 & \text{on } \partial\Omega \times (0, T) \\ u(x, 0) = u_0 & \text{in } \Omega \end{cases} \quad (P_\lambda)$$

where $\lambda \geq 0$ and $\chi > 0$ are a constant. In [19] F.H. Lin and Y. Yang by using variational argument derived the following nonlocal MEMS equation

$$\begin{cases} -\Delta v = \frac{\lambda}{(1-v)^2(1 + \chi \int_{\Omega} \frac{dx}{1-v})^2} & \text{in } \Omega \\ v = 0 & \text{on } \partial\Omega \end{cases}$$

for modeling the stationary deflection between the two parallel plates of an electrostatic MEMS device with circuit series capacitance.

Note that λ is proportional to the square of the voltage. When the voltage is very large, the two membranes will touch each other or quench and we would expect the

solutions of (P'_λ) and (P_λ) cease to exist after a finite time. One would like to get estimates for the touchdown time of the solutions of (P_λ) and (P'_λ) . We refer the readers to the papers [4], [5], [6], by N. Ghoussoub and Y. Guo for various touchdown time estimates for the solutions of (P'_λ) .

In this paper we will obtain touchdown time estimates for the solutions of (P_λ) . We prove that the quenching set of (P_λ) is compact under a mild condition on the initial value of the solution. When $\Omega = B_R = \{x \in \mathbb{R}^n : |x| < R\}$ and u_0 is radially symmetric and monotone decreasing in $0 \leq r \leq R$, we prove that the solution u of (P_λ) satisfies

$$1 - u(x, t) \geq C|x|^2 \quad \text{in } B_R \times [t_0, T] \quad (1)$$

for some constants $t_0 \in (0, T)$, $C > 0$, and the point $x = 0$ is the only possible quenching set. When $\Omega = B_R$ and u_0 is radially symmetric which also satisfies some strict concavity assumption, we prove that for any $\beta \in (2, 3)$ the solution u of (P_λ) satisfies

$$1 - u(x, t) \geq C|x|^{\frac{2}{\beta}} \quad \text{in } Q_R^T \quad (2)$$

for some constant $C > 0$ and that the solution u quenches in a finite time for any sufficiently large $\lambda > 0$. We also obtain the quenching time estimate in this case.

We will assume that $\Omega \subset \mathbb{R}^n$ is a bounded C^2 domain and $\lambda > 0$, $\chi \geq 0$, for the rest of the paper. We start with some definitions. For any $\delta > 0$, $R > 0$, $T > 0$, let $\Omega_\delta = \{x \in \Omega : \text{dist}(x, \partial\Omega) < \delta\}$ and $Q_R^T = B_R \times (0, T)$. Observe that when Ω is a bounded convex domain then there exists a constant $\delta_0 > 0$ such that for any $x \in \Omega_{\delta_0}$ there exists a unique point $y \in \partial\Omega$ such that the line segment $\overline{xy}$ is perpendicular to $\partial\Omega$ at y .

For any constants $\chi \geq 0$, $\lambda > 0$, and

$$u_0 \in L^1(\Omega) \text{ with } u_0 \leq a \text{ a.e. in } \Omega \quad (3)$$

for some constant $0 < a < 1$ we say that u is a solution (subsolution, supersolution respectively) of (P_λ) in $\Omega \times (0, T)$ if $u \in C^{2,1}(\Omega \times (0, T)) \cap C(\overline{\Omega} \times (0, T))$, $u < 1$, satisfies

$$\frac{\partial u}{\partial t} = \Delta u + \frac{\lambda}{(1 - u)^2(1 + \chi \int_\Omega \frac{dy}{1 - u(y, t)})^2} \quad \text{in } \Omega \times (0, T)$$

($\leq$, $\geq$ respectively) in the classical sense with $u(x, t) = 0$ ($\leq$, $\geq$ respectively) on $\partial\Omega \times (0, T)$,

$$\sup_{\overline{\Omega} \times [0, T']} u(x, t) < 1 \quad \forall 0 < T' < T,$$

and

$$\|u(\cdot, t) - u_0\|_{L^1(\Omega)} \rightarrow 0 \quad \text{as } t \rightarrow 0.$$

For any solution u of (P_λ) we define the quenching time or touchdown time $T_\lambda > 0$ as the time which satisfies

$$\begin{cases} \sup_{\Omega} u(x, t) < 1 & \forall 0 < t < T_\lambda \\ \lim_{t \nearrow T_\lambda} \sup_{\Omega} u(x, t) = 1. \end{cases}$$

We say that u has a finite quenching time if $T_\lambda < \infty$ and we say that u quenches at time infinity if $T_\lambda = \infty$. For any solution u of (P_λ) we let the quenching set of u to be the set of points $x \in \Omega$ such that there exists a sequence $(x_k, t_k) \in \Omega \times (0, T_\lambda)$ such that $x_k \rightarrow x$ and $u(x_k, t_k) \rightarrow 1$ as $k \rightarrow \infty$.

We first recall some results of [15] and [16].

Theorem 1. (Theorem 2.1 and Theorem 2.2 of [16]) Let

$$-b \leq u_0 \leq a \quad \text{in } \Omega \quad (4)$$

for some constants $0 < a < 1$ and $b \geq 0$. Then for any $\lambda > 0$ and $\chi > 0$ there exists $T > 0$ such that (P_λ) has a unique solution $-b \leq u < 1$ in $\Omega \times (0, T)$ which satisfies

$$u(x, t) = \int_{\Omega} G(x, y, t) u_0(y) dy + \lambda \int_0^t \int_{\Omega} \frac{G(x, y, t-s)}{(1-u(y, s))^2 (1 + \chi \int_{\Omega} \frac{dz}{1-u(z, s)})^2} dy ds \quad (5)$$

for all $(x, t) \in \Omega \times (0, T)$ where $G(x, y, t)$ is the Dirichlet Green function for the heat equation in $\Omega \times (0, T)$.

Lemma 2. (cf. Theorem 2.1 of [15]) Let $u_{0,1}, u_{0,2} \in L^1(\Omega)$ be such that $0 \leq u_{0,1} \leq u_{0,2} \leq a$ in Ω for some constant $0 < a < 1$. Let $\lambda > 0$ and $0 \leq f \in C(\overline{\Omega} \times (0, T)) \cap L^\infty(\overline{\Omega} \times (0, T))$. Suppose u_1, u_2 , are nonnegative subsolution and supersolution of (P'_λ) in $\Omega \times (0, T)$ with initial value $u_0 = u_{0,1}, u_{0,2}$, respectively. Then $u_1 \leq u_2$ in $\overline{\Omega} \times (0, T)$.

By an argument similar to the proof of Proposition 2.1 of [5] we have the following theorem.

Lemma 3. Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain. Let u_0 satisfy (4) for some constants $0 < a < 1$ and $b \geq 0$. Suppose there exists a constant $\delta_0 > 0$ such that

$$u_0|_{\overline{\Omega}_{\delta_0}} \in C^1(\overline{\Omega}_{\delta_0}) \quad \text{and} \quad \frac{\partial u_0}{\partial \nu} < 0 \quad \text{on } \partial\Omega \quad (6)$$

where $\partial/\partial \nu$ is differentiation with respect to the unit outward normal ν on $\partial\Omega$ and u is the unique solution of (P_λ) in $\Omega \times (0, T)$ given by Theorem 1. Then there exists a constant $\delta_1 \in (0, \delta_0)$ such that for any $x \in \overline{\Omega}_{\delta_1}$ if y is the unique point on $\partial\Omega$ such that the line segment $\overline{xy}$ is perpendicular to $\partial\Omega$ at y , then

$$\frac{\partial}{\partial \vec{n}} u(z, t) < 0 \quad \forall z \in \overline{xy}, 0 < t < T$$

where $\partial/\partial \vec{n}$ is differentiation with respect to the unit vector $\vec{n}$ along the direction $\overline{xy}$.

By Lemma 3 and an argument similar to the proof of Proposition 1.3 of [16] and Propostion 2.1 of [9] we have the following two corollaries.

Corollary 4. *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain. Let u_0 satisfy (4) and (6) for some constants $0 < a < 1$, $b \geq 0$, and $\delta_0 > 0$. Suppose u is the unique solution of (P_λ) in $\Omega \times (0, T)$ given by Theorem 1. Let δ_1 be as in Theorem 3. Then there exist $a_1 > 0$ and $\delta \in (0, \delta_1)$ such that for any $y \in \Omega_\delta$ there exists a fixed-sized cone $\Gamma(y) \subset \Omega_{2\delta}$ with vertex at y such that $|\Gamma(y) \setminus \Omega_\delta| \geq a_1$ and $u(z, t) \geq u(y, t)$ for any $z \in \Gamma(y)$ and $0 < t < T$. Moreover*

$$\int_{\Omega_\delta} \frac{dx}{(1 - u(x, t))^2} \leq \frac{|\Omega|}{a_1} \int_{\Omega \setminus \Omega_\delta} \frac{dx}{(1 - u(x, t))^2} \quad \forall 0 < t < T. \quad (7)$$

Corollary 5. *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain. Let u_0 satisfy (4) and (6) for some constants $0 < a < 1$, $b \geq 0$, and $\delta_0 > 0$. Suppose u is the unique solution of (P_λ) in $\Omega \times (0, T)$ given by Theorem 1 such that u touchdown at time T . Then the set of touchdown points for u is a compact subset of Ω .*

By (7) and an argument similar to the proof of Theorem 4.3 of [16] we have the following theorem.

Theorem 6. *Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain. Let u_0 satisfy (4) and (6) for some constants $0 < a < 1$, $b = 0$, and $\delta_0 > 0$. Let $\chi > 0$. Then there exists a constant $C_1 > 0$ and such that for any $\lambda > \lambda_0 = C_1 \mu_1$ and any solution u of (P_λ) , u quenches in a finite time*

$$T_\lambda \leq \frac{C_1}{\lambda - \lambda_0}.$$

Lemma 7. *Let $\lambda > 0$, $\chi > 0$, and $\Omega \subset \mathbb{R}^n$ be a bounded domain such that $B_R \subset \Omega$ for some constant $R > 0$. Let $0 \leq u_0 \leq a$ in Ω for some constant $0 < a < 1$ and let $0 \leq u < 1$ be a solution of (P_λ) in $\Omega \times (0, T)$ with*

$$\sup_{0 \leq t < T} \int_{\Omega} \frac{dz}{1 - u(z, t)} < \infty.$$

Let

$$A(t) = \left(1 + \chi \int_{\Omega} \frac{dz}{1 - u(z, t)} \right)^{-2} \quad (8)$$

and $0 < \delta_1 \leq \inf_{0 \leq t < T} A(t)$. Suppose $\lambda > 2n/\delta_1 R^2$. Then T satisfies

$$T \leq \frac{1}{\lambda \delta_1} \left(1 - \frac{2n}{\lambda \delta_1 R^2} \right)^{-1}. \quad (9)$$

Proof: Let $v = 1 - u$ and $v_0 = 1 - u_0$. Then v satisfies

$$\begin{cases} v_t = \Delta v - \lambda A(t)v^{-2} & \text{in } \Omega \times (0, T) \\ v(x, t) = 1 & \text{on } \partial\Omega \times (0, T) \\ v(x, 0) = v_0 & \text{in } \Omega. \end{cases} \quad (10)$$

Let

$$\psi(x, t) = 1 - \lambda\delta_1 c_0 t \left(1 - \frac{|x|^2}{R^2}\right)$$

where

$$c_0 = 1 - \frac{2n}{\lambda\delta_1 R^2}.$$

Then

$$\frac{2n}{R^2} \cdot \frac{1}{\lambda\delta_1 c_0} = c_0^{-1} - 1. \quad (11)$$

Hence by (11),

$$\psi_t - \Delta\psi + \lambda\delta_1\psi^{-2} \geq \lambda\delta_1 c_0 [-1 - (2nt/R^2) + c_0^{-1}] \geq 0 \quad \text{in } Q_R^{T_1}$$

where $T_1 = \min(T, 1/(\lambda\delta_1 c_0))$. Hence ψ is a supersolution of

$$\begin{cases} \psi_t = \Delta\psi - \lambda\delta_1\psi^{-2} & \text{in } Q_R^{T_1} \\ \psi = 1 & \text{on } \partial B_R \times [0, T_1) \cap \overline{B}_R \times \{0\}. \end{cases} \quad (12)$$

Suppose $T > 1/(\lambda\delta_1 c_0)$. Since v is a subsolution of (12), by Lemma 2,

$$\begin{aligned} 0 < v &\leq \psi \quad \text{in } \overline{Q_R^{T_1}} \\ \Rightarrow 0 < v(0, 1/(\lambda\delta_1 c_0)) &\leq \psi(0, 1/(\lambda\delta_1 c_0)) = 0. \end{aligned}$$

Contradiction arises and (9) follows. $\square$

Theorem 8. Let $0 \leq u_0 \leq a$ be a radially symmetric function in $B_R \subset \mathbb{R}^n$ for some constant $0 < a < 1$ which is monotone decreasing in $0 \leq r \leq R$. Let $T > 0$. Suppose u is the unique solution of (P_λ) in Q_R^T given by Theorem 1 and u touches down at time T . Then there exist constants $t_0 \in (0, T)$ and $C > 0$ depending on λ and χ such that (1) holds. Hence $x = 0$ is the only quenching point of u at time T . Moreover for $n \geq 3$ we have

$$\sup_{0 \leq t < T} \int_{B_R} \frac{dz}{1 - u(z, t)} < \infty. \quad (13)$$

Proof: We first observe that by an argument similar to the proof of Theorem 4.3 of [16] and Theorem 1.5 of [13] $1 > u(x, t) = u(|x|, t) \geq 0$ is radially symmetric in Q_R^T with

$$u_r(0, t) = 0 \quad \text{and } u_r(r, t) < 0 \quad \forall 0 < r \leq R, 0 < t < T. \quad (14)$$

Let

$$v_1(x, t) = \int_{B_R} G(x, y, t) u_0(y) dy$$

and

$$F(x, t, s) = \lambda \int_{B_R} G(x, y, t - s) A(s) (1 - u(y, s))^{-2} dy$$

where $A(t)$ is given by (8) with $\Omega = B_R$. Since $0 \leq v_1 \leq a$ satisfies the heat equation in Q_R^T with $v_1 \equiv 0$ on $\partial B_R \times (0, T)$ and u_0 is radially symmetric and monotone decreasing for $0 \leq r \leq R$, by the maximum principle and an argument similar to the proof of Theorem 1.5 of [13] v_1 is radially symmetric and $v_{1,r}(r, t) \leq 0$ for any $0 < r \leq R$ and $0 < t < T$.

Similarly for any $0 < s < T$, $F(\cdot, \cdot, s)$ is a radially symmetric solution of the heat equation in $B_R \times (s, T)$ with $F(x, t, s) \equiv 0$ for any $x \in \partial B_R$, $0 < s < t < T$, and

$$\frac{\partial F}{\partial r}(r, t, s) < 0 \quad \forall 0 < r = |x| \leq R, 0 < s < t < T. \quad (15)$$

By Theorem 1 u satisfies (5). By (5) and (15),

$$u_r(r, t) = v_{1,r}(r, t) + \int_0^t F_r(r, t, s) ds \leq \int_0^{\frac{T}{3}} F_r(r, t, s) ds \quad \forall 0 \leq r \leq R, T/3 \leq t < T. \quad (16)$$

Let

$$v_2(x, t) = \int_0^{\frac{T}{3}} F(x, t, s) ds \quad \forall |x| \leq R, T/3 \leq t < T. \quad (17)$$

Then v_2 is radially symmetric and satisfies

$$\begin{cases} v_{2,t} = \Delta v_2 = \frac{1}{r^{n-1}}(r^{n-1}v_{2,r})_r & \text{in } B_R \times (T/3, T) \\ v_2 = 0 & \text{on } \partial B_R \times (T/3, T) \\ v_2 > 0 & \text{in } B_R \times \{T/3\} \end{cases} \quad (18)$$

By (15) for each $T/3 < t < T$ $v_{2,r}(r, t)$ is monotone decreasing in $0 \leq r \leq R$. Hence for each $T/3 < t < T$ $v_2(x, t)$ will attain its maximum at $x = 0$. Differentiating (18) with respect to r and letting $q = r^{n-1}v_{2,r}$,

$$q_t = q_{rr} - \frac{n-1}{r}q_r \quad \text{in } (0, R) \times (T/3, T). \quad (19)$$

Let $\tilde{q} = q + \varepsilon r^n$ for some constant $\varepsilon > 0$ to be determined later. Then

$$\tilde{q}_t - \tilde{q}_{rr} + \frac{n-1}{r}\tilde{q}_r = 0 \quad \text{in } (0, R) \times (T/3, T) \quad (20)$$

and

$$\tilde{q}(0, t) = 0 \quad \forall T/3 < t < T. \quad (21)$$

By the maximum principle ([1], [18]) there exists a constant $C_1 > 0$ such that

$$v_{2,r}(R, t) \leq -C_1 \quad \forall T/2 \leq t < T.$$

Hence by choosing $0 < \varepsilon \leq \varepsilon_1 = C_1/(2R)$ we have

$$\tilde{q}(R, t) \leq -(C_1/2)R^{n-1} \quad \forall T/2 \leq t < T. \quad (22)$$

By the strong maximum principle,

$$\begin{aligned} v_2(r, t') &< \max_{y \in \overline{B}_R} v_2(y, t) = v_2(0, t) \quad \forall 0 \leq r \leq R, T/2 \leq t < t' < T \\ \Rightarrow v_2(0, t') &< v_2(0, t) \quad \forall T/2 \leq t < t' < T \\ \Rightarrow v_{2,t}(0, t) &\leq 0 \quad \forall T/2 \leq t < T. \end{aligned} \quad (23)$$

Suppose $v_{2,t}(0, t) \equiv 0$ on $(T/2, T)$. Then

$$v_2(0, t') = v_2(0, t) \quad \forall T/2 \leq t < t' < T.$$

This contradicts (23). Hence there exists $t_0 \in (T/2, T)$ such that

$$v_{2,t}(0, t_0) < 0. \quad (24)$$

By (18),

$$v_{2,t}(r, t_0) = v_{2,rr}(r, t_0) + \frac{n-1}{r}v_{2,r}(r, t_0). \quad (25)$$

Since $v_{2,r}(0, t) = 0$ for any $T/3 \leq t < T$, letting $r \rightarrow 0$ in (25) by the l'hospital rule,

$$v_{2,t}(0, t_0) = nv_{2,rr}(0, t_0). \quad (26)$$

By (24) and (26),

$$v_{2,rr}(0, t_0) < 0.$$

Then there exists $0 < r_1 < R$ such that

$$\varepsilon_2 = -\max_{0 \leq r \leq r_1} v_{2,rr}(r, t_0) > 0. \quad (27)$$

Let $\varepsilon_3 = -\max_{r_1 \leq r \leq R} v_{2,r}(r, t_0)$. Then $\varepsilon_3 > 0$. Let $\varepsilon_0 = \min(\varepsilon_1, \varepsilon_2/2, \varepsilon_3/(2R))$ and $0 < \varepsilon \leq \varepsilon_0$. Since $v_{2,r}(0, t_0) = 0$, by the mean value theorem for any $0 < r \leq r_1$ there exists $0 \leq r' \leq r$ such that

$$\tilde{q}(r, t_0) = r^n \left(\frac{v_{2,r}(r, t_0)}{r} + \varepsilon \right) = r^n (v_{2,rr}(r', t_0) + \varepsilon) < 0. \quad (28)$$

and

$$\tilde{q}(r, t_0) = r^{n-1} (v_{2,r}(r, t_0) + \varepsilon r) \leq r^{n-1} (-\varepsilon_3 + \varepsilon R) < 0 \quad \forall r_1 \leq r \leq R. \quad (29)$$

By (28) and (29),

$$\tilde{q}(r, t_0) < 0 \quad \forall 0 < r \leq R. \quad (30)$$

By (20), (21), (22), (30) and the strong maximum principle,

$$\begin{aligned} \tilde{q}(r, t) &< 0 \quad \forall 0 < r \leq R, t_0 \leq t < T \\ \Rightarrow v_{2,r}(r, t) &\leq -\varepsilon r \quad \forall 0 < r \leq R, t_0 \leq t < T \end{aligned} \quad (31)$$

Hence by (16) and (31),

$$u_r(r, t) \leq v_{2,r}(r, t) \leq -\varepsilon r \quad \forall 0 < r \leq R, t_0 \leq t < T.$$

Thus

$$u(r, t) = u(0, t) + \int_0^r u_r(\rho, t) d\rho \leq 1 - \varepsilon \int_0^r \rho d\rho \leq 1 - (\varepsilon/2)r^2$$

holds for any $0 \leq r \leq R$, $t_0 \leq t < T$, and (1) follows. By (1) $x = 0$ is the only quenching point of u and for any $n \geq 3$,

$$\int_{B_R} \frac{dz}{1 - u(z, s)} \leq C \int_0^R r^{n-3} dr \leq CR^{n-2} \quad \forall t_0 \leq s < T$$

and (13) follows. $\square$

Theorem 9. *Let $0 \leq u_0 \leq a$ be a radially symmetric C^2 function on $\overline{B}_R \subset \mathbb{R}^n$ for some constant $0 < a < 1$ which is monotone decreasing in $0 \leq r \leq R$ with*

$$u_0 \equiv 0 \quad \text{on } \partial B_R \quad \text{and} \quad u_0''(r) \leq -c_1 \quad \forall 0 \leq r \leq R \quad (32)$$

for some constant $c_1 > 0$. Let $\chi > 0$ and $T > 0$. Then there exists a constant $\lambda_0 > 0$ such that for any $\lambda \geq \lambda_0$ and $2 < \beta < 3$ if u is the unique solution of (P_λ) in Q_R^T given by Theorem 1, then there exists a constant $C > 0$ such that (2) holds and for any $n \in \mathbb{Z}^+$ (13) holds. Moreover there exists a constant $\lambda_1 \geq \lambda_0$ such that if $\lambda > \lambda_1$ and T is the maximal time of existence of the solution u of (P_λ) , then T satisfies (9) and $x = 0$ is the only quenching point of u at the quenching time T .

Proof. Note that the estimate (2) for the case $n = 1$ and sufficiently large λ is proved in [8]. Here we will modify the proof of [8] so that it works for any dimension n . Let $\lambda \geq \lambda_0$ and $2 < \beta < 3$ for some constant $\lambda_0 > 0$ to be determined later. Let u be the unique solution of (P_λ) in Q_R^T given by Theorem 1. By an argument similar to the proof of Theorem 8 $0 \leq u < 1$ is radially symmetric in Q_R^T and u satisfies (14). Let $v = 1 - u$ and $v_0 = 1 - u_0$. Then v satisfies (10). By (32),

$$v_0 = 1 \quad \text{on } \partial B_R \quad \text{and} \quad v_0'(r) \geq c_1 r \quad \forall 0 \leq r \leq R. \quad (33)$$

By (10), (33), and an argument similar to the proof of Proposition 2.1 of [9] and Theorem 4.1 of [8] there exists a constant $0 < c_2 < c_1$ depending on u_0 but is independent of u such that

$$v_r(R, t) \geq c_2 R \quad \forall 0 \leq t < T. \quad (34)$$

Let $w = v^\beta$. Then w satisfies

$$w_t - \frac{1}{r^{n-1}}(r^{n-1}w_r)_r = -\lambda\beta A(t)w^{1-\frac{3}{\beta}} - \frac{\beta-1}{\beta} \cdot \frac{w_r^2}{w} \quad \text{in } (0, R) \times (0, T) \quad (35)$$

with $w = 1$ on $\partial B_R \times (0, T)$. By (14) $w_r(r, t) \geq 0$ for any $0 \leq r \leq R$ and $0 \leq t < T$. Differentiating (35) with respect to r and setting $q = r^{n-1}w_r$, we have

$$\begin{aligned} q_t - q_{rr} + \frac{n-1}{r}q_r &= \lambda(3-\beta)A(t)w^{-\frac{3}{\beta}}q + \frac{\beta-1}{\beta}r^{n-1}\frac{w_r^3}{w^2} - 2\frac{(\beta-1)}{\beta}\frac{w_{rr}}{w}q \\ &\geq \lambda(3-\beta)A(t)w^{-\frac{3}{\beta}}q - 2\frac{(\beta-1)}{\beta}\frac{w_{rr}}{w}q \quad \text{in } (0, R) \times (0, T). \end{aligned} \quad (36)$$

Now

$$w_{rr}q = \left(\frac{q}{r^{n-1}}\right)_r q = \frac{q_r q}{r^{n-1}} - \frac{n-1}{r^n}q^2 \leq q_r w_r. \quad (37)$$

Since $0 < w \leq 1$, by (36) and (37),

$$\begin{aligned} q_t - q_{rr} + \frac{n-1}{r}q_r &\geq \frac{w_r}{w} \left(\lambda(3-\beta)A(t)r^{n-1}w^{1-\frac{3}{\beta}} - 2\frac{(\beta-1)}{\beta}q_r \right) \\ &\geq \frac{w_r}{w} \left(\lambda(3-\beta)A(t)r^{n-1} - 2\frac{(\beta-1)}{\beta}q_r \right) \quad \text{in } (0, R) \times (0, T). \end{aligned} \quad (38)$$

We now choose $\varepsilon > 0$ sufficiently small such that

$$A(0) > \delta_1 := [1 + \chi|\omega_{n-1}|(2/\varepsilon)^{\frac{1}{\beta}}(n - (2/\beta))^{-1}R^{n-\frac{2}{\beta}}]^{-2} \quad (39)$$

and

$$\varepsilon < \beta \min(c_1(1-a)^{\beta-1}, c_2) \quad (40)$$

where ω_{n-1} is the surface area of a unit ball in $\mathbb{R}^n$. We next choose

$$\lambda_0 > \frac{2\varepsilon n(\beta-1)}{\beta(3-\beta)\delta_1}.$$

Since $\lambda \geq \lambda_0$,

$$\lambda(3-\beta)\delta_1 > 2\frac{(\beta-1)}{\beta}\varepsilon n. \quad (41)$$

By (39) and (41) there exists $0 < T_1 \leq T$ such that $[0, T_1)$ is the maximal interval such that

$$\lambda(3 - \beta)A(t) > 2\frac{(\beta - 1)}{\beta}\varepsilon n \quad (42)$$

holds for any $0 \leq t < T_1$. Let $\tilde{q} = q - \varepsilon r^n$. Then by (38) and (42),

$$\tilde{q}_t - \tilde{q}_{rr} + \frac{n-1}{r}\tilde{q}_r + 2\frac{(\beta-1)}{\beta}\frac{w_r}{w}\tilde{q}_r \geq 0 \quad \text{in } (0, R) \times (0, T_1). \quad (43)$$

Now by (34) and (40),

$$\tilde{q}(R, t) = R^{n-1}(\beta v^{\beta-1}v_r(R, t) - \varepsilon R) = R^{n-1}(\beta v_r(R, t) - \varepsilon R) \geq 0 \quad \forall 0 \leq t < T. \quad (44)$$

By (33) and (40),

$$\tilde{q}(r, 0) \geq 0 \quad \forall 0 \leq r \leq R. \quad (45)$$

By (43), (44), (45) and the maximum principle,

$$\begin{aligned} & \tilde{q}(r, t) \geq 0 \quad \forall 0 \leq r \leq R, 0 \leq t < T_1 \\ \Rightarrow & w_r(r, t) \geq \varepsilon r \quad \forall 0 \leq r \leq R, 0 \leq t < T_1 \\ \Rightarrow & w(r, t) \geq w(0, t) + \frac{\varepsilon}{2}r^2 > \frac{\varepsilon}{2}r^2 \quad \forall 0 \leq r \leq R, 0 \leq t < T_1 \\ \Rightarrow & v(r, t) > \left(\frac{\varepsilon}{2}\right)^{\frac{1}{\beta}} r^{\frac{2}{\beta}} \quad \forall 0 \leq r \leq R, 0 \leq t < T_1. \end{aligned} \quad (46)$$

By (46),

$$\int_{B_R} \frac{dz}{v(z, t)} < |\omega_{n-1}|(2/\varepsilon)^{\frac{1}{\beta}}(n - (2/\beta))^{-1}R^{n-\frac{2}{\beta}} \quad \forall 0 \leq t < T_1 \quad (47)$$

$$\Rightarrow A(t) > \delta_1 \quad \forall 0 \leq t < T_1. \quad (48)$$

If $T_1 < T$, then by (41) and (48) there exists $T_2 \in (T_1, T)$ such that (42) holds on $(0, T_2)$. This contradicts the maximality of T_1 . Hence $T_1 = T$. By (46) and (47) we get that (2) and (13) hold.

We now let $\lambda > \lambda_1$ for some constant λ_1 to be determined later and let T be the maximal time of existence of the solution u of (P_λ) . Suppose $T > 1$. By repeating the above argument but with T being replaced by 1 in the argument and noting that then the constant c_2 is now independent of T we can find constants ε , δ_1 , and λ_0 as before which are then all independent of T . Then (2) holds in Q_R^1 and (13), (48), hold for any $0 \leq t \leq 1$. Let

$$\lambda_1 = \max(\lambda_0, 4n/(\delta_1 R^2), 3/\delta_1). \quad (49)$$

By (49) and Lemma 7,

$$1 \leq \frac{1}{\lambda\delta_1} \left(1 - \frac{2n}{\lambda\delta_1 R^2}\right)^{-1} \leq \frac{2}{\lambda\delta_1} \leq \frac{2}{3}.$$

Contradiction arises. Hence $T < 1$. Note that then we can choose the constant c_2 in the above argument to be independent of T . Hence the constants ε , δ_1 , and λ_0 are all independent of T . Then by (48) Lemma 7 holds. Hence T satisfies (9). Thus u quenches in a finite time T bounded above by the right hand side of (9). By (2) $x = 0$ is the only quenching point of u at the quenching time T and the theorem follows. $\square$

References

- [1] A. Friedman, *Partial differential equations of parabolic type*, Prentice-Hall, Inc., Englewood Cliffs, N.J., U.S.A., 1964.
- [2] G. Flores, G.A. Mercado and J.A. Pelesko, *Dynamics and Touchdown in Electrostatic MEMS*, Proceedings of ICMENS (2003), 182–187.
- [3] N. Ghoussoub and Y.J. Guo, *On the partial differential equations of electrostatic MEMS devices: stationary case*, *SIAM J. Math. Anal.* 38 (2007), no. 5, 1423–1449.
- [4] N. Ghoussoub and Y.J. Guo, *On the partial differential equations of electrostatic MEMS devices II: dynamic case*, *NoDEA Differential Equations Appl.* 15 (2008), no. 1-2, 115–145.
- [5] N. Ghoussoub and Y.J. Guo, *On the partial differential equations of electrostatic MEMS devices III: refined touchdown behavior*, *J. Diff. Eqns.* 244 (2008), no. 9, 2277–2309.
- [6] N. Ghoussoub and Y.J. Guo, *Estimates for the quenching time of a parabolic equation modeling electrostatic MEMS*, *Methods Appl. Anal.* 15 (2008), 361–376.
- [7] N. Ghoussoub and Y.J. Guo, *Pull-in voltage and steady states of nonlocal electrostatic MEMS*, preprint.
- [8] J.S. Guo, B. Hu and C.J. Wang, *A nonlocal quenching problem arising in a micro-electro mechanical system*, *Quart. Appl. Math.* 67 (2009), no. 4, 725–734.
- [9] Y.J. Guo, *On the partial differential equations of electrostatic MEMS devices III: Refined touchdown behavior*, *J. Diff. Eqns* 244 (2008), 2277–2309.
- [10] Y.J. Guo, Z.G. Pan and Michael J. Ward, *Touchdown and Pull-in Voltage Behavior of a MEMS Device with Varying Dielectric Properties*, *SIAM J. Appl. Math.* 66 (2005), no. 1, 309–338.
- [11] Z. Guo and J. Wei, *On the Cauchy problem for a reaction-diffusion equation with a singular nonlinearity*, *J. Diff. Eqns* 240 (2007), 279–323.

- [12] Z. Guo and J. Wei, *On a fourth order nonlinear elliptic equation with negative exponent*, SIAM J. Math. Anal. 40 (2008/09), no. 5, 2034–2054.
- [13] S.Y. Hsu, *Asymptotic behaviour of solutions of the equation $u_t = \Delta \log u$ near the extinction time*, Advances in Diff. Eqns. 8 (2003), no. 2, 161–187.
- [14] K.M. Hui, *Growth rate and extinction rate of a reaction diffusion equation with a singular nonlinearity*, Diff. and Int. Eqns 22 (2009), nos. 7–8, 771–786.
- [15] K.M. Hui, *Global and touchdown behaviour of the generalized MEMS device equation*, Adv. Math. Sci. Appl. 19 (2009), no. 1, 347–370.
- [16] K.M. Hui, *Existence and dynamic properties of a parabolic nonlocal MEMS equation*, <http://arxiv.org/abs/0809.4209v2>.
- [17] N.I. Kavallaris, T. Miyasita and T. Suzuki, *Touchdown and related problems in electrostatic MEMS device equation*, NoDEA Nonlinear Differential Equations Appl. 15 (2008), no. 3, 363–385.
- [18] O.A. Ladyzenskaya, V.A. Solonnikov, and N.N. Uraltceva, *Linear and quasilinear equations of parabolic type*, Transl. Math. Mono. Vol 23, Amer. Math. Soc., Providence, R.I., U.S.A., 1968.
- [19] F. Lin and Y. Yang, *Nonlinear non-local elliptic equation modelling electrostatic actuation*, Proc. Royal Soc. London, Ser. A 463 (2007), 1323–1337.
- [20] L. Ma and J.C. Wei, *Properties of positive solutions to an elliptic equation with negative exponent*, J. Functional Analysis 254 (2008), 1058–1087.
- [21] J.A. Pelesko, *Mathematical Modeling of Electrostatic MEMS with Tailored Dielectric Properties*, SIAM J. Appl. Math. 62 (2002), no. 3, 888–908.
- [22] J.A. Pelesko and D.H. Bernstein, *Modeling MEMS and NEMS*, Chapman Hall and CRC Press, 2002.
- [23] J.A. Pelesko and A.A. Triolo, *Nonlocal Problems in MEMS Device Control*, J. Eng. Math. 41 (2001), no. 4, 345–366.